WORK AND ENERGY EXERCISE 1

- If the unit of force and length be each increased by four times, then the unit of energy is increased by a) 16 times
 b) 8 times
 c) 2 times
 d) 4 times
- 2. A light and heavy body have equal momentum . Which one has greater kinetic energy ?a) the light bodyb) both have equal K.E.
 - c) the heavy body d) data given is incomplete
- 3. A light and heavy body have equal kinetic energy which has greater momentum ?a) the heavy bodyb) the light body
 - c) both have equal momentum d) data given is incomplte
- 4. A moving train is stopped by applying certain force. If the speed is doubled then the distance will be

IIT-JEE : PHYSICS : EXERCISE

	a) the same c) half	b) doubled d) four times	
5.	 A metal ball falls from a height of 10 metre on a steel plate and bounces back to a height of 2.5 metre The coefficient of restitution of the ball is 		
	a) 0.5	b) 0.25	
	c) 0.75	d) 0.33	
6.	. The kinetic energy acquired by a mass m in traveling a certain distance d, starting from rest, under the action of a constant force is directly proportional to		
	a) \sqrt{m}	b) independent of m	
	C) 1/ \sqrt{m}	d) m	
7.	7. A body moves a distance of 10 m along a straight line under the action of a 5 N force. If the workdone is 25 J, then angle between the force and direction of motion of the body is		
	a) 30°	b) 45°	
	c) 60°	d) 75°	
8.	 The kinetic energy acquired by a mass m after traveling a fixed distance from rest under the action of constant force is 		
	a) directly proportional to $\sqrt{\mathrm{m}}$	b) inversely proportional to $\sqrt{\mathrm{m}}$	
	c) directly proportional to m	d) independent of m	
9.	One man takes 1 minute to raise a box to a he the same. The energy of the two is	eight of 1 metre and another man takes 30 second to do	
	a) different	b) same	
	c) energy of the first is more	d) energy of the second is more	
10	10. A particle moves under the effect of a force $F = cx$ from $x = 0$ to $x = x_1$. The work done in the process is		
	a) cx_1^2	b) $\frac{1}{2}cx_1^2$	
	C) cx_1^3	d) zero	
11	 One kilowatt hour is equal to a) 36×10⁵ J 	b) 36×10 ³ J	
	C) 36×10 ^{−5} J	d) 36×10 ⁻³ J	

12. A body is moved along a straight line by a machine delivering constant power . The distance moved by		
the body in time t is proportional to		
a) t ^{1/2}	b) t ^{3/4}	

c) t ^{3/2}	d) t ²
$(1 + 3)^2$	d) +4
UIL	u) l

A uniform chain of length L and mass M is lying on a smooth table and one-third of its lying on a smooth table and one-third of its length is hanging vertically down over the edge of the table. If g is acceleration due to gravity, the work required to pull the hanging part on to the table is

 a) Mgl
 b) Mgl /3

a) MyL	b) NigL/ 5
c) MgL/9	d) mgL/18

14. A long spring is stretched by 2 cm , its potential energy is U . If the spring is stretched by 10 cm, the potential energy stored in it will be

a) U/25	b) U/5
c) 5 U	d) 25 U

15. A body moves a distance of 10 m along a straight line under the action of a force of 5 N. If the work done is 25 joule, the angle which the force makes with the direction of motion of the body is
a) 0°
b) 30°
c) 60°
d) 90°

EXERCISE II

A long spring is stretched by 2 cm. Its potential energy is U. If the spring is stretched by 10 cm, the potential energy stored in it will be

 a) 11/25
 b) 11/5

a) 0/25	D) U/5
c) 5 U	d) 25 U

2. A ball with kinetic energy E is thrown at an angle of 45° with horizonal , its kinetic energy at the highest point of its flight will be

a) E	b) E/ $\sqrt{2}$
c) E/2	d) zero

3. A bullet of mass a and velocity b is fired into a large block of mass c. The final velocity of the system is

a)
$$\left(\frac{c}{a+b}\right).b$$
 b) $\left(\frac{a}{a+c}\right).b$

C)
$$\left(\frac{a+b}{c}\right)a$$
 d) $\left(\frac{a+c}{a}\right)b$

4. A boy and a man carry a uniform rod of length L, horizontally in such a way that the boy gets 1/4th loads. If the boy is at one end of the rod, the distance of the man from the other end is

a) L/3	b) L/4
c) 2 L/3	d) 3 L/4

 If momentum of a certain body is increased by 50% then increase in the kinetic energy of the body will be

a) 25%	b) 50%
c) 100%	d) 125%

6. Two bodies of masses M_A and M_B move is opposite directions with velocities v_A and v_B respectively . After an elastic collision they exchange their velocities , then the ratio M_A/M_B must be

a) 1	b) 2
c) 4	d) 8/9

7. A position dependent force, $F = 7 - 2x + 3x^2$ newton acts on a small body of mass 2 kg and displaces it from x = 0 to x = 5 m. The workdone in joule is

- a) 70 b) 270 c) 35 d) 125
- 8. The potential energy between two atoms, in a molecule, is given by

$$U(x) = \frac{a}{x^{12}} - \frac{b}{x^6}$$

where a and b are positive constants and x is the distance between the atoms . The atom is in stable equilibrium, when

a) x = 0c) $x = (2a / b)^{1/6}$ b) $x = (a / 2b)^{1/6}$ d) $x = (11a/5b)^{1/6}$

- **9.** A body of mass m accelerates from rest to v_1 in time t_1 . As a function of time t, the instantaneous power delivered to the body is
 - a) $\frac{m_1 v_1}{t_1}$ b) $\frac{m v_1^2 t}{t_1}$ c) $\frac{m_1 v_1^2 t}{t_1^2}$ d) $\frac{m v_1 t^2}{t_1}$

- **10**. A car of mass m is driven with acceleration a long a straight level and road against a constant external resistive force R. When the velocity of car is v, the rate at which the engine of the car is doing work will be
 - a) Rv b) mav c) (R + ma) v d) (ma – R) v
- A body is moved along a straight line by a machine delivering constant power. The distance moved by the body in time t is proportional to
 - a) $t^{1/2}$ b) $t^{3/4}$ c) $t^{3/2}$ d) t^2
- 12. A running man has half the kinetic energy that a boy half his mass has. The man speeds up by 1 m/s and then has the same kinetic energy as the boy. The original speed of the man is

a) $(1+\sqrt{2}) m/s$	b) $(2+\sqrt{2})m/s$
C) $(3+\sqrt{2})m/s$	d) $\sqrt{2} m/s$

13. A wind –powered generator converts wind energy into electric energy. Assume that the generator coverts a fixed fraction of the wind energy intercepted by its blades into electric energy. For wind speed v, the electric power output will be proportional to

a) v	b) v ²
c) v ³	d) v ⁻¹

ANSWERS													
Exerc	ise I												
1a	2a	3a	4d	5a	6b	7c	8d	9b	10b				
11a	12c	13d	14d	15c									
Exerc	ise II												
1d	2c	3b	4b	5d	6a	7d	8c	9c	10c				
11c12a							13c						