

CHAPTER 11
DIFFERENTIAL EQUATIONS

TOPICS:

- 1. Differential Equation, Order And Degree**
- 2. Formation Of Differential Equation.**
- 3. Variable Separable Method**
- 4. Homogeneous Method**
- 5. Non Homogeneous Method**
- 6. Linear Differential Equation**
- 7. Bernoulli's Differential Equation**

DIFFERENTIAL EQUATIONS

An equation involving one dependent variable, one or more independent variables and the differential coefficients (derivatives) of dependent variable with respect to independent variables is called a differential equation.

ORDER OF A DIFFERENTIAL EQUATION :

The order of the highest derivative involved in an ordinary differential equation is called the order of the differential equation.

DEGREE OF A DIFFERENTIAL EQUATION

The degree of the highest derivative involved in an ordinary differential equation, when the equation has been expressed in the form of a polynomial in the highest derivative by eliminating radicals and fraction powers of the derivatives is called the degree of the differential equation.

EXERCISE --- 11(A)

I

- Find the order of the family of the differential equation obtained by eliminating the arbitrary constants b and c from $xy = ce^x + be^{-x} + x^2$.**

Sol.

Equation of the curve is $xy = ce^x + be^{-x} + x^2$

Number of arbitrary constants in the given curve is 2.

Therefore, the order of the corresponding differential equation is 2.

- Find the order of the differential equation of the family of all circles with their centers at the origin.**

Given family of curves is $x^2 + y^2 = a^2 \dots (1)$, a parameter.

Diff (1) w.r.t x, $2x+2y.y_1 = 0$.

Hence required differential equation is $x+y.y_1 = 0$.

Order of the differential equation is 1.

II

- Form the differential equation of the following family of curves where parameters are given in brackets.**

i). $y = c(x-c)^2; (c)$

$$y = c(x-c)^2 \dots (1)$$

Diff. w.r.t x,

$$y_1 = c.2(x-c) \dots (2)$$

$$\frac{(1)}{(2)} \Rightarrow \frac{y}{y_1} = \frac{x-c}{2}$$

$$\Rightarrow x-c = \frac{2y}{y_1} \quad \text{and} \quad c = x - \frac{2y}{y_1}$$

$$\text{from (1), } y = \left(x - \frac{2y}{y_1} \right) \left(\frac{2y}{y_1} \right)^2 \Rightarrow y^3 = 4y(xy_1 - 2y)$$

ii) $xy = ae^x + be^{-x}; (a, b)$

$$xy = ae^x + be^{-x} \dots (1)$$

Diff.w.r.t. x,

$$y + x.y_1 = ae^x - be^{-x} \dots (2)$$

diff.w.r.t. x,

$$y_1 + y_1 + xy_2 = ae^x + be^{-x} = xy$$

$$\therefore 2y_1 + xy_2 = xy$$

Which is required differential equation.

iii) $y = (a+bx)e^{kx}; (a, b)$

$$y = (a+bx)e^{kx} \dots (1)$$

Diff.w.r.t x,

$$\Rightarrow y_1 = k(a+bx)e^{kx} + be^{kx}$$

$$\Rightarrow y_1 = ky + be^{kx} \dots (2)$$

Diff.w.r.t. x,

$$\Rightarrow y_2 = ky_1 + kbe^{kx}$$

$$\Rightarrow y_2 = ky_1 + k(y_1 - ky)$$

$$\Rightarrow y_2 = 2ky_1 - k^2 y \quad \text{which is required differential equation.}$$

iv) $y = a \cos(nx+b); (a, b)$

ans: $y_2 + n^2 y = 0$

- 2. Obtain the differential equation which corresponds to each of the following family of curves.**

- i) The rectangular hyperbolas which have the coordinates axes as asymptotes.**

Sol. Equation of the rectangular hyperbola is $xy=c^2$ where c is arbitrary constant.

Differentiating w.r.t. x

$$x \frac{dy}{dx} + y = 0$$

ii) The ellipses with centres at the origin and having coordinate axes as axes.

Sol. Equation of ellipse is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

Diff. w.r.t.x,

$$\frac{2x}{a^2} + \frac{2y}{b^2} \frac{dy}{dx} = 0 \Rightarrow y \cdot y_1 = -\frac{b^2}{a^2} x$$

Diff. w.r.t. x,

$$y \cdot y_2 + y_1 \cdot y_1 = -\frac{b^2}{a^2} \Rightarrow y \cdot y_2 + 2y_1 = \frac{y \cdot y_1}{x}$$

$$\Rightarrow x(y \cdot y_2 + 2y_1) = y \cdot y_1$$

III.

1. Form the differential equations of the following family of curves where parameters are given in brackets.

i) $y = ae^{3x} + be^{4x}; (a, b)$

Sol. $y = ae^{3x} + be^{4x} \dots\dots(1)$

Differentiating w.r.t x

$$y_1 = 3ae^{3x} + 4be^{4x} \dots\dots(2)$$

Differentiating w.r.t x,

$$y_2 = 9ae^{3x} + 16be^{4x} \dots\dots(3)$$

Eliminating a,b from above equations,

$$\begin{vmatrix} y & e^{3x} & e^{4x} \\ y_1 & 3e^{3x} & 4e^{4x} \\ y_2 & 9e^{3x} & 16e^{4x} \end{vmatrix} = 0 \Rightarrow \begin{vmatrix} y & 1 & 1 \\ y_1 & 3 & 4 \\ y_2 & 9 & 16 \end{vmatrix} = 0$$

$\Rightarrow y_2 - 7y_1 + 12y = 0$ which is the required differential equation.

ii) $y = ax^2 + bx ; (a, b)$

Ans: $x^2 \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} + 2y = 0$

iii) $ax^2 + by^2 = 1$; (a, b)

Sol.

Given equation is

$$ax^2 + by^2 = 1 \dots\dots(1)$$

Differentiating w.r.t. x

$$\Rightarrow 2ax + 2byy_1 = 0$$

$$\Rightarrow ax + byy_1 = 0 \dots\dots(2)$$

Differentiating w.r.t. x

$$\Rightarrow a + b(yy_2 + y_1y_1) = 0 \Rightarrow a + b(yy_2 + y_1^2) = 0$$

$$\Rightarrow ax + bx(yy_2 + y_1^2) = 0 \dots\dots(3)$$

$$(3) - (2) \Rightarrow bx(yy_2 + y_1^2) - byy_1 = 0$$

$$\Rightarrow x(yy_2 + y_1^2) - yy_1 = 0$$

iv) $xy = ax^2 + \frac{b}{x}$; (a, b)

Ans: $x^2 \left(\frac{d^2y}{dx^2} \right) + 2x \left(\frac{dy}{dx} \right) - 2y = 0$

2. Obtain the differential equation which corresponds to each of the following family of curves.

i) The circles which touch the Y-axis at the origin.

Sol. Equation of the given family of circles is

$$x^2 + y^2 + 2gx = 0, g \text{ is arbitrary const } \dots(i)$$

$$x^2 + y^2 = -2gx$$

Differentiating w.r.t. x

$$2x + 2yy_1 = -2g \dots(ii)$$

Substituting in (i)

$$x^2 + y^2 = x(2x + 2yy_1) \text{ by (ii)}$$

$$= 2x^2 + 2xyy_1$$

$$yy^2 - 2xyy_1 - 2x^2 = 0$$

$$y^2 - x^2 = 2xy \frac{dy}{dx}.$$

- ii) The parabolas each of which has a latus rectum $4a$ and whose axes are parallel to x-axis.

Sol.

Equation of the given family of parabolas is

$$(y - k)^2 = 4a(x - h) \text{-----(i)}$$

where h, k are arbitrary constants

Differentiating w.r.t. x

$$2(y - k)y_1 = 4a$$

$$(y - k)y_1 = 2a \text{ ...(2)}$$

Differentiating w.r.t. x

$$(y - k)y_2 + y_1^2 = 0 \text{ ...(3)}$$

$$\text{From (2), } y - k = \frac{2a}{y_1}$$

Substituting in (3)

$$\frac{2a}{y_1} \cdot y_2 + y_1^2 = 0 \Rightarrow 2ay_2 + y_1^3 = 0$$

- iii) The parabolas having their foci at the origin and axis along the x-axis.

Sol.

Given family of parabolas is $y^2 = 4a(x + a)$ -----(i)

Diff. w.r.t. x ,

$$2y \frac{dy}{dx} = 4a \Rightarrow \frac{1}{2}yy' = a \text{ -----(2)}$$

From (i) and (2),

$$y^2 = 4 \frac{1}{2}yy' \left(x + \frac{1}{2}yy' \right)$$

$$y^2 = 2y'x + 4 \cdot \frac{1}{4}y^2y'^2 \Rightarrow y^2 = 2yy'x + y^2y'^2$$

$$y \left(\frac{dy}{dx} \right)^2 + 2x \left(\frac{dy}{dx} \right) = y$$

SOLUTIONS OF DIFFERENTIAL EQUATIONS

1. VARIABLES SEPARABLE

Let the given equation be $\frac{dy}{dx} = f(x, y)$. If $f(x, y)$ is a variables separable function, i.e., $f(x, y) = g(x)h(y)$ then the equation can be written as $\frac{dy}{dx} = g(x)h(y) \Rightarrow \frac{dy}{h(y)} = g(x)dx$. By integrating both sides, we get the solution of $\frac{dy}{dx} = f(x, y)$. This method of finding the solution is known as variables separable.

EXERCISE – 11(B)

1. Find the general solution of $\sqrt{1-x^2}dy + \sqrt{1-y^2}dx = 0$.

Sol. Given D.E is $\sqrt{1-x^2}dy + \sqrt{1-y^2}dx = 0$

$$\sqrt{1-x^2}dy = -\sqrt{1-y^2}dx$$

Integrating both sides

$$\int \frac{dy}{\sqrt{1-y^2}} = - \int \frac{dx}{\sqrt{1-x^2}}$$

$$\sin^{-1}y = -\sin^{-1}x + c$$

Solution is $\sin^{-1}x + \sin^{-1}y = c$, where c is a constant.

2. Find the general solution of $\frac{dy}{dx} = \frac{2y}{x}$.

Sol. $\frac{dy}{dx} = \frac{2y}{x} \Rightarrow \int \frac{dy}{y} = 2 \int \frac{dx}{x}$

Integrating both sides

$$\log c + \log y = 2 \log x$$

$$\log cy = \log x^2$$

Solution is $cy = x^2$ where c is a constant.

II. Solve the following differential equations.

1. $\frac{dy}{dx} = \frac{1+y^2}{1+x^2}$

Sol. $\frac{dy}{dx} = \frac{1+y^2}{1+x^2}$

Integrating both sides

$$\Rightarrow \int \frac{dy}{1+y^2} = \int \frac{dx}{1+x^2}$$

$$\Rightarrow \tan^{-1} y = \tan^{-1} x + \tan^{-1} c \text{ where } c \text{ is a constant.}$$

2. $\frac{dy}{dx} = e^{y-x}$

Sol. $\frac{dy}{dx} = \frac{e^y}{e^x} \Rightarrow \frac{dy}{e^y} = \frac{dx}{e^x}$

Integrating both sides $\int e^{-x} dx = \int e^{-y} dy \Rightarrow -e^{-x} = -e^{-y} + c$

$$e^{-y} = e^{-x} + c \text{ where } c \text{ is a constant.}$$

3. $(e^x + 1)y dy + (y + 1)dx = 0$

Sol. $(e^x + 1)y dy = -(y + 1)dx$

$$\frac{ydy}{y+1} = -\frac{dx}{e^x + 1}$$

Integrating both sides

$$\int \left(1 - \frac{1}{y+1}\right) dy = \int -\frac{e^{-x} dx}{e^{-x} + 1}$$

$$y - \log(y+1) = \log(e^{-x} + 1) + \log c$$

$$\Rightarrow y - \log(y+1) = \log c(e^{-x} + 1)$$

$$\Rightarrow y = \log(y+1) + \log c(e^{-x} + 1)$$

$$y = \log c(y+1)(e^{-x} + 1)$$

Solution is : $e^y = c(y+1)(e^{-x} + 1)$.

4. $\frac{dy}{dx} = e^{x-y} + x^2 e^{-y}$

Sol. $\frac{dy}{dx} = e^{x-y} + x^2 e^{-y} = \frac{e^x}{e^y} + \frac{x^2}{e^y}$

Integrating both sides

$$\int e^y \cdot dy = \int (e^x + x^2) dx$$

Solution is : $e^y = e^x + \frac{x^3}{3} + c$

$$5. \tan y \, dx + \tan x \, dy = 0$$

Sol. $\tan y \, dx = -\tan x \, dy$

$$\frac{dx}{\tan x} = \frac{-dy}{\tan y} \Rightarrow \frac{\cos x}{\sin x} dx = -\frac{\cos y}{\sin y} dy$$

Taking integration

$$\int \frac{\cos x}{\sin x} dx = - \int \frac{\cos y}{\sin y} dy$$

$$\log \sin x = -\log \sin y + \log c$$

$$\log \sin x + \log \sin y = \log c$$

$$\log(\sin x \cdot \sin y) = \log c \Rightarrow \sin x \cdot \sin y = c$$

$$6. \sqrt{1+x^2} dx + \sqrt{1+y^2} dy = 0$$

$$\text{Sol. } \sqrt{1+x^2} dx = -\sqrt{1+y^2} dy$$

$$\text{Integrating both sides } \int \sqrt{1+x^2} dx = - \int \sqrt{1+y^2} dy$$

$$\frac{x}{2} \times \sqrt{1+x^2} + \frac{1}{2} \sinh^{-1} x =$$

$$y \frac{\sqrt{1+y^2}}{2} = \frac{1}{2} \sinh^{-1} x + c$$

$$x\sqrt{1+x^2} + y\sqrt{1+y^2} + \log \left[(x+\sqrt{1+x^2})(y+\sqrt{1+y^2}) \right] = c$$

$$7. \quad y - x \frac{dy}{dx} = 5 \left(y^2 + \frac{dy}{dx} \right)$$

$$\text{Sol. } y - 5y^2 = (x+5) \frac{dy}{dx} \Rightarrow \frac{dx}{x+5} = \frac{dy}{y(1-5y)}$$

Integrating both sides

$$\int \frac{dx}{x+5} = \int \frac{dy}{y(1-5y)} = \int \left(\frac{1}{y} + \frac{5}{1-5y} \right) dy$$

$$\ln|x+5| = \ln y - \ln|1-5y| + \ln c$$

$$\ln|x+5| = \ln \left| \frac{cy}{1-5y} \right| \Rightarrow x+5 = \left(\frac{cy}{1-5y} \right)$$

8. $\frac{dy}{dx} = \frac{xy + y}{xy + x}$

Sol. $\frac{dy}{dx} = \frac{y(x+1)}{x(y+1)} \Rightarrow \frac{y+1}{y} dy = \frac{x+1}{x} dx$

$$\int \left(1 + \frac{1}{y}\right) dy = \int \left(1 + \frac{1}{x}\right) dx$$

$$y + \log y = x + \log x + \log c$$

$$y - x = \log \left| \frac{cx}{y} \right|$$

III.

1. $\frac{dy}{dx} = \frac{1+y^2}{(1+x^2)xy}$

Sol. $\frac{dy}{dx} = \frac{1+y^2}{(1+x^2)xy}$

$$\Rightarrow \frac{ydy}{1+y^2} = \frac{dx}{x(1+x^2)}$$

$$\frac{2ydy}{1+y^2} = \frac{2xdx}{x^2(1+x^2)}$$

Integrating both sides

$$\int \frac{2ydy}{1+y^2} = \int \left(\frac{1}{x^2} - \frac{1}{1+x^2} \right) 2x dx$$

$$\log(1+y^2) = \log x^2 - \log(1+x^2) + \log c$$

$$\log(1+x^2) + \log(1+y^2) = \log x^2 + \log c$$

$$\text{Solution is : } (1+x^2)(1+y^2) = cx^2$$

2. $\frac{dy}{dx} + x^2 = x^2 \cdot e^{3y}$

Sol. $\frac{dy}{dx} + x^2 = x^2 \cdot e^{3y}$

$$\Rightarrow \frac{dy}{dx} = x^2 \cdot e^{3y} - x^2 = x^2(e^{3y} - 1)$$

Integrating both sides

$$\int \frac{dy}{e^{3y} - 1} = \int x^2 dx \Rightarrow \int \frac{e^{-3y}}{1-e^{-3y}} = \int x^2 dx$$

$$\log \frac{(1-e^{-3y})}{3} = \frac{x^3}{3} + c$$

$$\log(1-e^{-3y}) = x^3 + c' \quad (c' = 3c)$$

$$\text{Solution is: } 1-e^{-3y} = e^{x^3} \cdot k \quad (k = e^{c'})$$

3. $(xy^2 + x)dx + (yx^2 + y)dy = 0$

Sol. $(xy^2 + x)dx + (yx^2 + y)dy = 0$

$$x(y^2 + 1)dx + y(x^2 + 1)dy = 0$$

$$\text{Dividing with } (1+x^2)(1+y^2)$$

$$\frac{x dx}{1+x^2} + \frac{y dy}{1+y^2} = 0$$

Integrating both sides

$$\int \frac{x dx}{1+x^2} + \int \frac{y dy}{1+y^2} = 0$$

$$\frac{1}{2} \left[(\log(1+x^2) + \log(1+y^2)) \right] = \log c$$

$$\log(1+x^2)(1+y^2) = 2 \log c = \log c^2$$

$$(1+x^2)(1+y^2) = k \text{ when } k = c^2.$$

4. $\frac{dy}{dx} = 2y \tanh x$

Sol. $\frac{dy}{dx} = 2y \tanh x \Rightarrow \frac{dy}{y} = 2 \tanh x dx$

Integrating both sides

$$\int \frac{dy}{y} = 2 \int \tanh x dx$$

$$\log y = 2 \log |\cosh x| + \log c$$

$$\ln y = 2 \ln \cosh x + \ln c \Rightarrow y = c \cosh^2 x$$

5. $\sin^{-1} \left(\frac{dy}{dx} \right) = x + y$

$$\frac{dy}{dx} = \sin(x+y) \Rightarrow x+y = t$$

$$1 + \frac{dy}{dx} = \frac{dt}{dx}$$

$$\frac{dt}{dx} - 1 = \sin t \Rightarrow \frac{dt}{dx} = 1 + \sin t$$

$$\frac{dt}{1+\sin t} = dx$$

Integrating both sides

$$\int \frac{dt}{1+\sin t} = \int dx$$

$$\int \frac{1-\sin t}{\cos^2 t} dt = x + c$$

$$\int \sec^2 t dt - \int \tan t \cdot \sec t dt = x + c$$

$$\tan t - \sec t = x + c$$

$$\Rightarrow \tan(x+y) - \sec(x+y) = x + c$$

$$6. \quad \frac{dy}{dx} + \frac{y^2 + y + 1}{x^2 + x + 1} = 0$$

$$\frac{-dy}{y^2 + y + 1} = \frac{dx}{x^2 + x + 1}$$

Integrating both sides

$$-\int \frac{dy}{y^2 + y + 1} = \int \frac{dx}{x^2 + x + 1}$$

$$-\int \frac{dy}{\left(y + \frac{1}{2}\right)^2 + \frac{3}{4}} = \int \frac{dx}{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}}$$

$$-\frac{2}{\sqrt{3}} \tan^{-1} \frac{(y+1/2)}{\sqrt{3/2}} = \frac{2}{\sqrt{3}} \tan^{-1} \frac{(x+1/2)}{\sqrt{3/2}} + c$$

$$\tan^{-1} \frac{2x+1}{\sqrt{3}} + \tan^{-1} \frac{2y+1}{\sqrt{3}} = c$$

$$7. \quad \frac{dy}{dx} = \tan^2(x+y)$$

$$\text{Sol. } \frac{dy}{dx} = \tan^2(x+y) \text{ put } v = x+y$$

$$\frac{dv}{dx} = 1 + \frac{dy}{dx} = 1 + \tan^2 v = \sec^2 v$$

$$\int \frac{dv}{\sec^2 v} = \int dx = \int \cos^2 v \cdot dv = x + c$$

$$\int \frac{(1+\cos 2v)}{2} dv = x + c$$

$$\Rightarrow \int (1+\cos 2v) dv = 2x + 2c$$

$$\begin{aligned}
 v + \frac{\sin 2v}{2} &= 2x + 2c \\
 2v + \sin 2v &= 4x + c' \\
 2(x+y) + \sin 2(x+y) &= 4x + c' \\
 x - y - \frac{1}{2} \sin[2(x+y)] &= c
 \end{aligned}$$

Homogeneous Differential Equations :

A differential equation $\frac{dy}{dx} = \frac{f(x,y)}{g(x,y)}$ is said to be a homogeneous differential equation in x, y if

both $f(x, y)$, $g(x, y)$ are homogeneous functions of same degree in x and y.

To find the solution of the h.d.e put $y = vx$, then $\frac{dy}{dx} = v + x \frac{dv}{dx}$. Substituting these values in given differential equation, then it reduces to variable separable form. Then we find the solution of the D.E.

NOTE: Some times to solve the give homogeneous differential equation, we take the substitution $x = vy$.

EXERCISE – 11(C)

- 1. Express $x dy - y dx = \sqrt{x^2 + y^2} dx$ in the form $F\left(\frac{y}{x}\right) = \frac{dy}{dx}$.**

Sol. $x \cdot dy - y dx = \sqrt{x^2 + y^2} dx$

$$x \frac{dy}{dx} - y = \sqrt{x^2 + y^2} \Rightarrow x \frac{dy}{dx} = y + \sqrt{x^2 + y^2}$$

$$\frac{dy}{dx} = \frac{y}{x} + \sqrt{\frac{x^2 + y^2}{x^2}} = \frac{y}{x} + \sqrt{1 + \left(\frac{y}{x}\right)^2}$$

Which is of the form $F\left(\frac{y}{x}\right) = \frac{dy}{dx}$

- 2. Express $\left(x - y \tan^{-1} \frac{y}{x}\right) dx + x \tan^{-1} \frac{y}{x} dy = 0$ in the form $F\left(\frac{y}{x}\right) = \frac{dy}{dx}$.**

Ans: $\frac{dy}{dx} = \frac{\frac{y}{x} \cdot \tan^{-1}\left(\frac{y}{x}\right) - 1}{\tan^{-1}\left(\frac{y}{x}\right)} = F\left(\frac{y}{x}\right)$

3. Express $x \cdot \frac{dy}{dx} = y(\log y - \log x + 1)$ in the form $F\left(\frac{y}{x}\right) = \frac{dy}{dx}$.

$$\text{Ans: } \frac{dy}{dx} = \frac{y}{x} \left(\log \frac{y}{x} + 1 \right)$$

II. Solve the following differential equations.

$$1. \quad \frac{dy}{dx} = \frac{x-y}{x+y}$$

$$\text{Sol. } \frac{dy}{dx} = \frac{x-y}{x+y} \quad \dots \dots (1)$$

(1) is a homogeneous D.E.

Put $y = vx$

$$\frac{dy}{dx} = v + x \frac{dv}{dx}$$

$$v + x \frac{dv}{dx} = \frac{x - vx}{x + vx} = \frac{x(1-v)}{x(1+v)}$$

$$x \cdot \frac{dv}{dx} = \frac{1-v}{1+v} - v = \frac{1-v-v-v^2}{1+v} = \frac{1-2v-v^2}{1+v}$$

$$\int \frac{(1+v)dv}{1-2v-v^2} = \int \frac{dx}{x}$$

$$-\frac{1}{2} \log(1-2v-v^2) = \log x + \log c$$

$$-\frac{1}{2} \log \left(1 - 2 \cdot \frac{y}{x} - \frac{y^2}{x^2} \right) = \log cx$$

$$\log \frac{(x^2 - 2xy - y^2)}{x^2} = -2 \log cx = \log(cx)^{-2}$$

$$\frac{x^2 - 2xy - y^2}{x^2} = (cx)^{-2} = \frac{1}{c^2 x^2}$$

$$(x^2 - 2xy - y^2) = \frac{1}{c^2} = k$$

$$2. \quad (x^2 + y^2)dy = 2xy \, dx$$

Sol. $\frac{dy}{dx} = \frac{2xy}{x^2 + y^2}$ which is a homogeneous D.E.

Put $y = vx$

$$\frac{dy}{dx} = v + x \cdot \frac{dv}{dx}$$

$$v + x \cdot \frac{dv}{dx} = \frac{2x(vx)}{x^2 + v^2 x^2} = \frac{2v}{1+v^2}$$

$$x \cdot \frac{dv}{dx} = \frac{2v}{1+v^2} - v = \frac{2v-v-v^3}{1+v^2} = \frac{v-v^3}{1+v^2}$$

$$\int \frac{1+v^2}{v(1-v^2)} dv = \int \frac{dx}{x}$$

$$\text{Let } \frac{1+v^2}{v(1-v^2)} = \frac{A}{v} + \frac{B}{1+v} + \frac{C}{1-v}$$

$$1+v^2 = A(1-v^2) + BV(1-v) + CV(1+v)$$

$$v=0 \Rightarrow 1=A$$

$$v=1 \Rightarrow 1+1=C(2) \Rightarrow C=1$$

$$v=-1 \Rightarrow 1+1=B(-1)(2) \Rightarrow 2=-2B \Rightarrow B=-1$$

$$\int \frac{1+v^2}{v(1-v^2)} dv = \int \frac{dv}{v} - \int \frac{dv}{1+v} + \int \frac{dv}{1-v}$$

$$= \log v - \log(1+v) - \log(1-v) = \log \frac{v}{1-v^2}$$

$$\therefore \log \frac{v}{1-v^2} = \log x + \log c = \log cx$$

$$\frac{v}{1-v^2} = cx \Rightarrow v = cx(1-v^2)$$

$$\frac{y}{x} = cx \left(1 - \frac{y^2}{x^2}\right) \Rightarrow \frac{y}{x} = cx \frac{(x^2 - y^2)}{x^2}$$

Solution is: $y = c(x^2 - y^2)$

$$3. \quad \frac{dy}{dx} = \frac{-(x^2 + 3y^2)}{(3x^2 + y^2)}$$

Sol. $\frac{dy}{dx} = \frac{-(x^2 + 3y^2)}{(3x^2 + y^2)}$ which is a homogeneous D.E.

Put $y = vx$

$$\frac{dy}{dx} = v + x \cdot \frac{dv}{dx}$$

$$v + x \cdot \frac{dv}{dx} = \frac{-(x^2 + 3v^2 x^2)}{3x^2 + v^2 x^2} = \frac{-x^2(1+3v^2)}{x^2(3+v^2)}$$

$$x \cdot \frac{dv}{dx} = -v - \frac{1+3v^2}{3+v^2}$$

$$= \frac{-3v - v^3 - 1 - 3v^2}{3+v^2} = -\frac{(v+1)^3}{3+v^2}$$

$$\frac{3+v^2}{(v+1)^3} = \frac{-dx}{x}$$

$$\frac{3+v^2}{(v+1)^3} = \frac{A}{v+1} + \frac{B}{(v+1)^2} + \frac{C}{(v+1)^3}$$

Multiplying with $(v+1)^3$

$$3+v^2 = A(v+1)^2 + B(v+1) + C$$

$$v = -1 \Rightarrow 3+1 = C \Rightarrow C = 4$$

Equating the coefficients of v^2

$$A = 1$$

Equating the coefficients of V

$$0 = 2A + B$$

$$B = -2A = -2$$

$$\frac{v^2+3}{(v+1)^3} = \frac{1}{v+1} - \frac{2}{(v+1)^2} + \frac{4}{(v+1)^3}$$

$$\int \frac{v^2+3}{(v+1)^3} dx = - \int \frac{dx}{x}$$

$$\int \left(\frac{1}{v+1} - \frac{2}{(v+1)^2} + \frac{4}{(v+1)^3} \right) dv = -\log x + \log c \log(v+1) + \frac{2}{v+1} - \frac{4}{2(v+1)^2} = \log \frac{c}{x}$$

Solution is:

$$\log\left(\frac{y}{x}+1\right) + \frac{2}{\frac{y}{x}+1} - \frac{2}{\left(\frac{y}{x}+1\right)^2} = \log \frac{c}{x}$$

$$\frac{2x}{x+y} - \frac{2x^2}{(x+y)^2} = \log \frac{c}{x} - \log \frac{(x+y)}{x}$$

$$\frac{2x^2 + 2xy - 2x^2}{(x+y)^2} = \log \frac{c}{x+y}$$

$$\frac{2xy}{(x+y)^2} = \log \frac{c}{x+y}$$

$$\log\left(\frac{x+y}{c}\right)c = -\log\left(\frac{c}{x+y}\right) = -\frac{2xy}{(x+y)^2}$$

4. $y^2 dx + (x^2 - xy)dy = 0$

Sol. $y^2 dx + (x^2 - xy)dy = (xy - x^2)dy$

$$\frac{dy}{dx} = \frac{y^2}{xy - x^2} \text{ which is a homogeneous D.E.}$$

$$\text{Let } y = vx \Rightarrow \frac{dy}{dx} = v + x \cdot \frac{dv}{dx}$$

$$v + x \frac{dv}{dx} = \frac{v^2 x^2}{x^2(v-v^2)}$$

$$x \cdot \frac{dv}{dx} = \frac{v^2}{v-1} - v = \frac{v^2 - v^2 + v}{v-1}$$

$$\frac{v-1}{v} dv = \frac{dx}{x} \Rightarrow \int \left(1 - \frac{1}{v}\right) dv = \int \frac{dx}{x}$$

$$v - \log v = \log x + \log k$$

$$v = \log v + \log x + \log k = \log k(vx)$$

$$\frac{y}{x} = \log ky \Rightarrow ky = e^{y/x}$$

5. $\frac{dy}{dx} = \frac{(x+y)^2}{2x^2}$

Sol. $\frac{dy}{dx} = \frac{(x+y)^2}{2x^2}$ which is a homogeneous D.E.

$$y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$$

$$v + x \frac{dv}{dx} = \frac{(x + vx)^2}{2x^2} = x^2 \frac{(1+v)^2}{2x^2}$$

$$x \frac{dv}{dx} = \frac{(1+v^2)}{2} - v = \frac{1+v^2+2v-2v}{2}$$

$$2 \int \frac{dv}{1+v^2} = \int \frac{dx}{x} \Rightarrow 2 \tan^{-1} v = \log x + \log c$$

$$2 \tan^{-1} \left(\frac{y}{x} \right) = \log cx$$

6. $(x^2 - y^2)dx - xy dy = 0$

Ans: $x^2(x^2 - 2y^2) = k$

7. $(x^2y - 2xy^2)dx = (x^3 - 3x^2y)dy$

Sol. $(x^2y - 2xy^2)dx = (x^3 - 3x^2y)dy$

$$\frac{dy}{dx} = \frac{x^2y - 2xy^2}{x^3 - 3x^2y} \text{ which is a homogeneous D.E.}$$

Put $y = vx$ so that $\frac{dy}{dx} = v + x \frac{dv}{dx}$

$$\begin{aligned} v + x \frac{dv}{dx} &= \frac{x^3v - 2v^2x^3}{x^3 - 3vx^3} \\ &= \frac{(v - 2v^2)x^3}{(1 - 3v)x^3} = \frac{v - 2v^2}{1 - 3v} \end{aligned}$$

$$\begin{aligned} x \frac{dv}{dx} &= \frac{v - 2v^2}{1 - 3v} - v \\ &= \frac{v - 2v^2 - v(1 - 3v)}{1 - 3v} = \frac{-2v^2 + 3v^2}{1 - 3v} \end{aligned}$$

$$x \frac{dv}{dx} = \frac{v^2}{1 - 3v} \Rightarrow \frac{1 - 3v}{v^2} dv = \frac{dx}{x}$$

$$\int \left(\frac{1}{v^2} - \frac{3}{v} \right) dv = \int \frac{dx}{x}$$

$$\frac{-1}{v} - 3 \log v = \log x + \log c$$

$$\frac{-x}{y} = 3 \log \left(\frac{y}{x} \right) = \log x + \log c$$

$$\frac{-x}{y} - \log \left(\frac{y}{x} \right)^3 = \log xc$$

$$\frac{-x}{y} = \log xc + \log \frac{y^3}{x^3}$$

$$\frac{-x}{y} = \log \left(cx \cdot \frac{y^3}{x^3} \right) = \log \left(\frac{cy^3}{x^2} \right)$$

$$\frac{cy^3}{x^2} = e^{-x/y} \Rightarrow cy^3 = \frac{x^2}{e^{x/y}}$$

$$cy^3 \cdot e^{x/y} = x^2$$

8. $y^2 dx + (x^2 - xy + y^2)dy = 0$

Ans: $y = c \cdot e^{\tan^{-1}(y/x)}$

9. $(y^2 - 2xy)dx + (2xy - x^2)dy = 0$

Ans: $xy(y - x) = c$

10. $\frac{dy}{dx} + \frac{y}{x} = \frac{y^2}{x^2}$

Sol. $\frac{dy}{dx} + \frac{y}{x} = \frac{y^2}{x^2}$ which is a homogeneous D.E.

$$\text{Put } y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$$

$$v + x \frac{dv}{dx} + v = \frac{v^2 x^2}{x^2} \Rightarrow x \frac{dv}{dx} = v^2 - 2v$$

$$\frac{dv}{v^2 - 2v} = \frac{dy}{x}$$

$$\text{Let } \frac{1}{v^2 - 2v} = \frac{A}{v} + \frac{B}{v-2}$$

$$1 = A(v-2) + Bv$$

$$v=0 \Rightarrow 1=A(-2) \Rightarrow -\frac{1}{2}$$

$$v=2 \Rightarrow 1=2B \Rightarrow B=\frac{1}{2}$$

$$-\frac{1}{2} \int \left(\frac{1}{v} - \frac{1}{v-2} \right) dv = \int \frac{dx}{x}$$

$$-\frac{1}{2} [\log v - \log(v-2)] = \log x + \log c$$

$$-\frac{1}{2} \left[\log \frac{v}{v-2} \right] = \log cx$$

$$\log \frac{v}{v-2} = -\log cx = \log(cx)^{-2}$$

$$\frac{v}{v-2} = (cx)^{-2} \Rightarrow \frac{(y/x)}{(y/x)-2} = \frac{1}{c^2 x^2}$$

$$\frac{y}{y-2x} = \frac{1}{c^2 x^2} \Rightarrow x^2 y = \frac{1}{c^2} (y-2x)$$

Solution is :

$$y-2x = c^2 x^2 y = kx^2 y \text{ where } k=c^2$$

$$11. \quad xdy - ydx = \sqrt{x^2 + y^2} dx$$

$$\text{Ans: } y + \sqrt{x^2 + y^2} = cx^2$$

$$12. \quad (2x - y)dy = (2y - x)dx$$

$$\text{Ans; } (y-x) = c^2 (x+y)^3.$$

$$13. \quad (x^2 - y^2) \frac{dy}{dx} = xy$$

$$\text{Ans: } x^2 + 2y^2 (c + \log y) = 0.$$

$$14. \quad \text{Solve } 2 \frac{dy}{dx} = \frac{y}{x} + \frac{y^2}{x^2}$$

$$\text{Ans: } (y-x) = cx^2 y$$

III.

1. Solve $(1+e^{x/y})dx + e^{x/y} \left(1 - \frac{x}{y}\right)dy = 0$.

Sol. $(1+e^{x/y})dx + e^{x/y} \left(1 - \frac{x}{y}\right)dy = 0$

$$\Rightarrow \frac{dx}{dy} = -\frac{e^{x/y} \left(1 - \frac{x}{y}\right)}{(1+e^{x/y})} \text{ which is a homogeneous D.E.}$$

Put $x = vy \Rightarrow \frac{dx}{dy} = v + y \frac{dv}{dy}$

$$(1+e^v) \frac{dx}{dy} + e^v (1-v) = 0$$

$$(1+e^v) \left(v + y \frac{dv}{dy}\right) + e^v (1-v) = 0$$

$$v + ve^v + y(1+e^v) \frac{dv}{dy} + e^v - ve^v = 0$$

$$y(1+e^v)dv = -(v+e^v)dy$$

$$\int \frac{1+e^v}{v+e^v} dv = - \int \frac{dy}{y}$$

$$\log(v+e^v) = -\log y + \log c \Rightarrow v+e^v = \frac{c}{y}$$

$$\frac{x}{y} + e^{x/y} = \frac{c}{y} \Rightarrow x + y \cdot e^{x/y} = c$$

2. Solve : $x \sin \frac{y}{x} \cdot \frac{dy}{dx} = y \sin \frac{y}{x} - x$

Sol. $x \sin \frac{y}{x} \cdot \frac{dy}{dx} = y \sin \frac{y}{x} - x$

$$\Rightarrow \frac{dy}{dx} = \frac{\frac{y}{x} \left(\sin \left(\frac{y}{x} \right) - \frac{x}{y} \right)}{\sin \left(\frac{y}{x} \right)}$$

Put $y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$

$$v + x \frac{dv}{dx} = \frac{v \left(\sin v - \frac{1}{v} \right)}{\sin v}$$

$$x \frac{dv}{dx} = \frac{v \sin v - 1 - v \sin v}{\sin v}$$

$$-\sin v dv = \frac{1}{x} dx$$

$$\Rightarrow \int -\sin v \cdot dv = + \int \frac{dx}{x}$$

$$\Rightarrow \cos v = \log x + \log c = \log cx$$

$$\Rightarrow cx = e^{\cos v} = e^{\cos(y/x)}.$$

3. Solve: $x dy = \left(y + x \cos^2 \frac{y}{x} \right) dx$.

Ans: $\tan \left(\frac{y}{x} \right) = \log x + c$.

4. Solve : $(x - y \log y + y \log x)dx + x(\log y - \log x)dy = 0$.

Ans: $= (x - y) \log x + y \log y$

5. Solve $(ydx + xdy)x \cos \frac{y}{x} = (xdy - ydx)y \sin \frac{y}{x}$

Ans: $xy \cos \left(\frac{y}{x} \right) = c$

6. Find the equation of a curve whose gradient is $\frac{dy}{dx} = \frac{y}{x} - \cos^2 \frac{y}{x}$, where $x > 0, y > 0$ and which passes through the point $(1, \pi/4)$.

Sol. $\frac{dy}{dx} = \frac{y}{x} - \cos^2 \frac{y}{x}$ which is homogeneous differential equation.

Put $y = vx$

$$\frac{dy}{dx} = v + x \frac{dv}{dx}$$

$$v + x \frac{dv}{dx} = v - \cos^2 v \Rightarrow \int \frac{dv}{\cos^2 v} = - \int \frac{dx}{x}$$

$$\int \sec^2 v = - \int \frac{dx}{x} \Rightarrow \tan v = - \log |x| + c$$

This curve passes through $(1, \pi/4)$

$$\tan \left(\frac{\pi}{4} \right) = c - \log 1 \Rightarrow c = 1$$

Equation of the curve is:

$$\tan v = 1 - \log |x| \Rightarrow \tan\left(\frac{y}{x}\right) = 1 - \log |x|$$

Equations Reducible to Homogeneous Form -Non Homogeneous Differential Equations

The differential equation of the form $\frac{dy}{dx} = \frac{ax + by + c}{a'x + b'y + c'}$ is called non homogeneous differential equation.